## **Description of Map Units**

Lava flows of the Broken Top area, Craters of the Moon lava field Units arranged by age; youngest unit located at top of lefthand column and become progressively older as move down toward right

## MAP SYMBOLS

Buffalo Caves: lava tube formed by the withdrawal of molten lava after the formation of a surface crust

Hiking trail Road (Surfaced)

**Contour line** 

Contact

Dashed where approximately located; relative ages of flows along contact indicated by Y, younger, O, older

Approximate fault line; cube on downthrown side

Scarp

Gravity fault: vertical lines point to area of lower elevation

Eruptive and noneruptive fissures: eruptive fissures are fractures through which lava erupted, while noneruptive fissures represent fractures in the lava



**Kipuka** 

Area of older lava surrounded by younger lava flows; older lava was a topographic high when it was enclosed by younger flows. A kipuka is generally covered in mature vegetation and is a Hawaiian term for "island"



fa1

Broken Top flow (nd) and related deposits Pahoehoe basalt flow

through which lava did not erupt


Surface-fed flow from vents, fissures, cracks, and a few small spatter vents on east and southeast flanks of Broken Top cinder cone (ca1). Vents may have formed during partial collapse of the cone. Flow is characterized by hummocky, rough lava with polygonal joints, numerous inflation pits, pressuse plateaus, and ridges




Pressure plateau
Pressure induced uplift of lava flow, usually followed by a breakout flow and displaying numerous tension



Areas of pahoehoe where the surface of which contains open tubes and blisters







¢a1

fa2

fa2

fa2

ca1

Volcanic ash deposits

Ash and spatter from vent on west side of Broken Top cinder cone, generally deposited less than 1 m thick





Area on west-northwest side of Broken Top cinder cone characterized by numerous fault lines highlighting slump



Blue Dragon flow (2,076±45) and related deposits Pahoehoe basalt-hawaiite flow

Characterized by spiny pahoehoe flows having fresh, irridescent dark and light blue, glassy, vesicular surfaces. The flow consists of two lobes that moved east and west from fissure-controlled vents at the Spatter Cones (ca2) and the southern part of Big Craters cinder cone (ca5). The fissure system features spatter cones and pit craters.



Formed near rootless vents from degassed, highly viscous lava that was erupted late in the eruptive sequence, generally over relatively steep slopes





Lava onlap related to inflation from lava ponds



(Big Sink and Unnamed Sink)













The Broken Top Area was first mapped by Mel Kuntz et al. (1989) as part of the U.S Geological Survey's 1:24,000 Geologic Map of the Inferno Cone Quadrangle, Butte County, Idaho. More detailed mapping of the Broken Top Area was undertaken during the summer of 2004 by GIP (Geologist in the Parks) interns and other volunteers (Clennon, Truitt, and Hansen). This mapping was then completed during the summer of 2005 by a GIP intern (Wetherell). The purpose of the project was fourfold: first, the field mapping provided young geologists with experience in the field and allowed them to hone their previously acquired field mapping skills; second, the completed map provides Craters of the Moon National Monument and Preserve with an educational and interpretive tool for the public; third, the map offers a more detailed look at the youngest lava flow and associated cinder cone in the park; and finally, the Broken Top Area is important geologically because it exhibits nearly every feature associated with basaltic volcanism.

## Flow front

Related to Sawtooth flow; two prominent flow fronts ranging in height and length, surrounded by smaller fronts. Area is alsocharacterized by rafted blocks ranging in size from < 1 m to ~ 20 x 30 m



Crater

Crater rim on volcanic vent: hachures point toward central depression



Lava pond surrounded by levees; hachures point away from ponded area



Trench Mortar Flat flow (2,205±25) and related deposits Pahoehoe basalt-hawaiite flows fa3

Thin, shelly flows erupted (and surface fed) from the northern part of the Trench Mortar Flat eruptive fissures lying southeast of Big Cinder Butte cinder cone (cc3)



Accumulations of agglutinated spatter and cinders related to the Trench Mortar Flat events



Volcanic ash deposits aa3

Ash and fine lapilli associated with the Trench Mortar Flat eruptive events. The material is mainly < 1 m thick and is located along the north end of Trench Mortar Flat on the downwind (NE) side of the source vents



Sawtooth flow (6,020±160) and related deposits A'a trachyandesite flow

Flow is generally characterized by large ash units and steep-fronted lobes and was issued from vent at south end of Big Cinder Butte cinder cone (cc3)



Slabby lava flow fc3

fc3

cc3


fh1


Rafted slabs and surfaces of pahoehoe lava believed to occur as late-stage streams between Sawtooth A'a lobes. Some of the slabby flows appear to be remobilized regions of the Sawtooth A'a flow.



Big Cinder Butte cinder cone

Contains large crater with walls mantled by agglutinated spatter and minor flows. Reported by Stearns (1928) to be one of the largest purely basaltic cinder cones in the world, having a height of 240 m and a volume of approximately 0.2 cubic km. The aligned craters on the east and south flanks of the cone are likely to be part of the Trench Mortar





Ash and fine lapilli associated with Big Cinder Butte (cc3). Deposits occur as thick as 1 m on inner crater walls, and as thick as ~ 5 m, though discontinuously, on west and south flanks of cone



Crescent Butte flow (nd, ~15,000) and related deposits Pahoehoe basalt flow

Surface-fed, pahoehoe flow issued from Crescent Butte cinder cone (ch1). Majority of flow vegetated and covered in thin deposit of colluvium



Crescent Butte cinder cone ch1

Western side of cone is covered in ash from Trench Mortar Flat eruptive fissures. Cinder cone is the source vent for the Crescent Butte lava flow (fh1)



Cinder cones (Holocene and latest Pleistocene)

A series of Holocene and latest Pleistocene cinder cones having no known associated lava flows; exact ages have not been assigned to these cinder cones



Data for the 1:12,000 Broken Top Area map was drawn from U.S. Geological Survey publication GQ-1632 (Kuntz et al, 1989) and additional field mapping conducted during the summer of 2004 and 2005. GPS coordinates were recorded using a Trimble GeoXM GPS (Global Positioning System) unit and were transferred into ArcGIS 8.3 and 9.0. Radiocarbon ages were drawn from U.S. Geological Survey publication GQ-1632 (Kuntz et al, 1989); for further description see Kuntz, Champion, and others (1986), Kuntz and others (1982), and Kuntz, Spiker, and others (1986). \* Ages in years before present, nd, not dated

KEFERENCE, Kuntz, M.A., Lefebvre, R. H., Champion, D. E., Skipp, B. A., 1989, Geologic Map of the Inferno Cone Quadrangle, Butte County, Idaho: U.S. Geological Survey Quadrangle Map GQ-1632, scale 1:24,000.
Margaret, G., McAfee Jr., R., Wolf, C. L., eds., 1972, Glossary of Geology: American Geological Institute.
Stearns, H. T., 1928, Craters of the Moon National Monument, Idaho: Idaho Bureau of Mines and Geology Bulletin 13, 57 p. Field photos courtesy of: Douglas Owen, Kimberly Truitt, Nichole Hansen, Bob Lillie, and Prudence Wetherell